



Published in final edited form as:

Life Sci. 2010 August 28; 87(9-10): 269–274. doi:10.1016/j.lfs.2010.06.013.

Oxygen and oxygenation in stem-cell therapy for myocardial infarction

Mahmood Khan¹, Pawel Kwiatkowski², Brian K. Rivera¹, and Periannan Kuppasamy¹

¹Davis Heart and Lung Research Institute, Division of Cardiovascular Medicine, Department of Internal Medicine, The Ohio State University, Columbus, OH 43210, USA

²Davis Heart and Lung Research Institute, Division of Cardiothoracic Surgery, Department of Internal Medicine, The Ohio State University, Columbus, OH 43210, USA

Abstract

Myocardial infarction (MI) is caused by deprivation of oxygen and nutrients to the cardiac tissue due to blockade of coronary artery. It is a major contributor to chronic heart disease, a leading cause of mortality in the modern world. Oxygen is required to meet the constant energy demands for heart contractility, and also plays an important role in the regulation of heart function. However, reoxygenation of the ischemic myocardium upon restoration of blood flow may lead to further injury. Controlled oxygen delivery during reperfusion has been advocated to prevent this consequence. Monitoring the myocardial oxygen concentration would play a vital role in understanding the pathological changes in the ischemic heart following myocardial infarction. During the last two decades, several new techniques have become available to monitor myocardial oxygen concentration *in vivo*. Electron paramagnetic resonance (EPR) oximetry would appear to be the most promising and reliable of these techniques. EPR utilizes crystalline probes which yield a single sharp line, the width of which is highly sensitive to oxygen tension. Decreased oxygen tension results in a sharpening of the EPR spectrum, while an increase results in widening. In our recent studies, we have used EPR oximetry as a valuable tool to monitor myocardial oxygenation for several applications like ischemia-reperfusion injury, stem-cell therapy and hyperbaric oxygen therapy. The results obtained from these studies have demonstrated the importance of tissue oxygen in the application of stem cell therapy to treat ischemic heart tissues. These results have been summarized in this review article.

Keywords

myocardial infarction; stem cell therapy; oxygenation; electron paramagnetic resonance; oximetry; mesenchymal stem cells; hyperbaric oxygenation

Introduction

Oxygen is hailed as “...the elixir of life – a wonder tonic, a cure for ageing, a beauty treatment and potent medical therapy” (Lane 2002). Oxygen plays a principal role in aerobic respiration. In aerobic organisms, cellular respiration involves enzyme-catalyzed oxidation

Address for correspondence: Periannan Kuppasamy, PhD, 420 W. 12th Avenue, Room 114, Columbus, OH 43210, kuppasamy.1@osu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

of fuel substrates, primarily by oxygen, to yield the energy required for biological processes. The physiological homeostasis of these organisms is strictly maintained by optimal cellular and tissue oxygenation status through complex oxygen-sensing mechanisms, signaling cascades, and transport processes. In the event of fluctuating oxygen levels leading to either an increase (hyperoxia) or decrease (hypoxia) in cellular oxygen, the organism faces a crisis involving depletion of energy reserves, altered cell-signaling cascades, oxidative reactions/events, and cell death or tissue damage. Particularly, hypoxia can lead to serious disorders such as ischemic stroke or myocardial infarction (MI). For example, during myocardial ischemia which occurs when a region of the heart is deprived of blood supply due to a blockage of a coronary artery, the region becomes oxygen-limited leading to myocardial cell death. If left untreated, MI will lead to congestive heart failure, which remains a major cause of morbidity and mortality in the world.

Despite the recent advances in reperfusion strategies and medical treatment, MI and its consequences remain a serious clinical concern. The prolonged imbalance between the supply and demand for oxygen and nutrients by the myocardial tissue leads to a sub-acute or chronic state of myocardial ischemia, resulting in a new state of equilibrium. Chronic ischemia contributes to poor quality of life in many patients suffering from coronary heart disease (Thom et al. 2006) and is an increasingly important aspect of clinical practice. In chronically ischemic myocardium, early revascularization may interrupt a vicious cycle of cardiomyocyte degradation that would otherwise lead to progressive fibrosis and/or contractile dysfunction.

Stem-cell therapy for treating myocardial infarction

The use of cell therapy to promote myocardial repair has gained profound scientific and public interest. The first study on cell therapy using intracoronary mononuclear bone-marrow cells (mBMC) after MI in humans was reported in 2002 (Strauer et al. 2002). Even though several randomized, controlled trials are using the same method of cell delivery (Martin-Rendon et al. 2008a; Martin-Rendon et al. 2008b), the primary endpoint has generally been associated with increase in left-ventricular (LV) ejection fraction (EF) after 3 to 6 months. Feasibility of this treatment has been confirmed, and it appears to be safe in the short term. However, the efficacy of this approach varies between studies. The possible explanations for these differences could be related to limited study samples, selection of measurement techniques and disparity in timing of applied treatment. Moreover, number of cells used for treatment along with selection of placebo treatment or protocols for cell processing and expansion could greatly contribute to differences in clinical outcome of performed studies (Arnesen et al. 2007; Egeland and Brinchmann 2007a; Egeland and Brinchmann 2007b; Seeger et al. 2007; van Beem et al. 2008).

Studies have shown that the grafting of skeletal myoblasts, fetal cardiomyocytes, or embryonic or bone marrow-derived stem cells into myocardial scar tissue has resulted in marginal improvements in cardiac function and in the attenuation of abnormal cardiac remodeling (Agbulut et al. 2004; Chiu et al. 1995; Ghostine et al. 2002; Kamihata et al. 2001; Orlic et al. 2001a; Orlic et al. 2001b; Pagani et al. 2003; Retuerto et al. 2004; Tse et al. 2003). The reason for such modest improvements was attributed to the limited survival of the transplanted cells in the infarcted myocardium (McConnell et al. 2005). The hypovascular nature of the infarcted tissue may severely compromise the availability of oxygen, nutrients, and growth factors essential for the survival, engraftment and differentiation of the transplanted cells. The local hypoxic environment in the infarcted myocardium might be the main impediment to the survival of the transplanted cells. However, it is not clear whether the oxygen concentration in the ischemic myocardium (infarcted area) is altered by strategies leading to regional angiogenesis and/or by cell

transplantation. It is also unknown whether there is a direct relationship between local oxygen concentration and transplanted cell survival.

Role of oxygen in stem-cell therapy

Oxygen tension plays an important role in the growth of stem cells in culture and significantly influences their expansion and differentiation (Ezashi et al. 2005; Ma et al. 2009; Salim et al. 2004; Wang et al. 2005). In response to acute hypoxia, cardiomyocytes have been shown to exhibit adaptations that may facilitate cell survival and develop tolerance to subsequent acute severe hypoxia (Silverman et al. 1997). However, the levels and roles of oxygen concentration at the sites of transplantation in the heart have not been investigated. This is primarily due to the significant technical challenges in obtaining reliable and repeated measurements of oxygen tension in a functional, beating organ during and after stem cell therapy.

A number of methods are available to obtain measurements of tissue oxygen concentration *in vivo* (Heidt et al. 2009; Mik et al. 2009; Swartz and Clarkson 1998; Vogt et al. 2009). Of these, electron paramagnetic resonance (EPR) oximetry would appear to be the most promising to obtain reliable, accurate, and repeated measurements over time (Ahmad and Kuppusamy). Our group used EPR oximetry to measure myocardial pO₂ in rodent (murine and rat) hearts under a variety of pathophysiological and therapeutic conditions, including ischemia, reperfusion, pharmacological intervention, and cell therapy. The results have shown significant variations and importance of oxygen tension during cardiac damage and repair

Most recent studies on stem-cell therapy for MI utilizing mesenchymal stem cells (MSCs) suggest that understanding the importance of the tissue micro-environment and how it may be manipulated is critical to realize the effective therapeutic potential of these cells (Chen et al. 2007; Djouad et al. 2007). Most recent study from our laboratory had shown that oxygen concentration is one of the vital components within the micro-environment (Chacko et al. 2009). Oxygen plays a significant role in the control and regulation of many physiological, metabolic, and signaling pathways involved in cellular engraftment and host-tissue regeneration (Chacko et al. 2009). It is also known that low oxygen tension is involved in holding the stem cells in a quiescent state in which they retain their plasticity (D'Ippolito et al. 2006).

A study by Rochefort et al has shown that MSCs are regularly observed in the circulating blood of rats and that the circulating pool of MSCs is consistently and substantially increased (15-fold) in animals exposed to chronic hypoxia (Rochefort et al. 2006). In several *in vitro* studies, low oxygen concentrations have been shown to stimulate the differentiation of MSCs into adipocyte-like phenotype (Fink et al. 2004; Lennon et al. 2001; Ren et al. 2006). Other researchers have reported suppressive effects of reduced oxygen tension on MSC plasticity (Fehrer et al. 2007; Potier et al. 2007). In this article we have summarized our findings that demonstrate the dynamics of oxygen concentration during pathophysiological changes leading to MI as well as modification of regional cardiac tissue oxygenation during cell therapy.

EPR Oximetry for monitoring myocardial oxygenation

Although several methods are used to measure oxygen concentration, a suitable technique for noninvasive, repeated, and reliable measurements of oxygen in the same tissue over time was previously not available. The methods which are commonly available to measure tissue oxygen concentration include: Clarke electrode, NADH fluorescence, phosphorescence-quenching, myoglobin saturation, and ¹H-NMR. EPR spectroscopy, a technique commonly

used for direct detection of free radicals and paramagnetic species, has recently been adapted to reliably and accurately quantify the concentration of molecular oxygen in viable cells and tissues (Ahmad and Kuppusamy 2010; Swartz and Clarkson 1998). EPR oximetry refers to the measurement of the partial pressure of oxygen (pO_2) by EPR spectroscopy. The principle of EPR oximetry is based on the paramagnetic characteristics of molecular oxygen, which in its ground state has two unpaired electrons, and undergoes spin exchange interaction with the paramagnetic EPR spin probe. This process is sensitive to oxygen content, with the relaxation rate of the spin probe increasing as a function of oxygen content (concentration/pressure). This increased spin-spin relaxation rate results in increased line broadening that is directly proportional to the oxygen content (Pandian et al. 2003).

EPR oximetry requires the implantation of an oxygen-sensing paramagnetic spin probe into the tissue of interest. The particulate probes sense and report pO_2 in the tissue *milieu*. The crystalline form of lithium octa-*n*-butoxy-substituted naphthalocyanine radical (LiNc-BuO) is a particulate oximetry spin probe that we have recently synthesized and validated for *in vivo* oximetry (Chacko et al. 2009; Pandian et al. 2003; Wisel et al. 2007). The LiNc-BuO crystals are composed of stacks of neutral radicals of lithiated naphthalocyanine macrocycles (Pandian et al. 2006). The EPR spectra of these particulates are characterized by a single and very narrow absorption peak due to the strong exchange coupling between the unpaired electrons within the molecular stack. The peak-to-peak linewidth of the EPR spectrum obtained is used to calculate pO_2 using a standard calibration curve (Pandian et al. 2003). The probe, in the form of submicron-sized (270 ± 120 nm) crystals (hereafter referred to as OxySpin), can be internalized in cells without compromising their function (Chacko et al. 2009; Khan et al. 2007; Wisel et al. 2007). Unique advantages of OxySpin probes are that they are retained in cells/tissues for substantially long periods of time, and do not alter the differentiation capability of the stem cells (Chacko et al. 2009), thus enabling continuous monitoring of myocardial pO_2 *in vivo* for weeks after stem cell transplantation in mouse heart. Figure 1 shows some typical EPR spectra obtained from murine hearts receiving OxySpin-labeled stem-cell transplants. The results demonstrated that high quality spectral data could be obtained from beating mouse hearts after several days or weeks after transplantation.

Despite being a unique and powerful method for noninvasive monitoring of oxygen concentration in the heart, EPR oximetry has certain limitations. A major limitation is that the method, at present, is applicable only to measurements in the hearts of small animals such as mice and rats. In larger animals, motional (contractile/respiratory) artifacts and loss of sensitivity due to increased depth from the chest wall are major concerns. Further advances in the EPR instrumentation are required to perform the measurements in large animals, including humans.

Myocardial oxygenation under conditions of prolonged ischemia (PI) and stem-cell treatment

The myocardial pO_2 measurements were performed using an *in vivo* EPR spectrometer (Magnetech, Berlin, Germany) equipped with automatic coupling and tuning controls for measurements in beating hearts. Microcrystals of LiNc-BuO were used as a probe for EPR oximetry. Mice and rats, under inhalation anesthesia (air mixed with 1.5-2% isoflurane), were implanted with OxySpin probes in the left-ventricular mid-myocardium. We have previously reported that the implantation of OxySpin probe is non-toxic to the heart and long-term myocardial pO_2 measurements could be performed in the same animal for up to 3 months (Figure 2) and possibly longer (Khan et al. 2009a). Hence, EPR oximetry has several significant advantages over other techniques for *in vivo* applications.

In our earlier reports, we have shown that the ischemic region of a mouse heart was significantly hypoxic for 4 wks after the induction of myocardial ischemia by permanent LAD ligation (Khan et al. 2007). Furthermore, there was no significant change in tissue pO_2 during the development of LV remodeling. Transplantation of skeletal myoblast (SM) cells in the ischemic region resulted in a significant increase in pO_2 compared to untreated ischemic tissue (Figure 3). Therefore, it is inferred that the transplanted SM cells may be responsible for the augmented myocardial pO_2 in the infarcted heart. Overall, we observed an increase of ~ 2 mmHg, which corresponds to an increase of ~ 2.5 μM of steady-state concentration of oxygen at the site of stem-cell therapy. Thus the magnitude of increase in oxygenation, *albeit* small, may still be sufficient for the survival of cells in the infarcted region (Khan et al. 2007). Therefore, it is imperative to investigate newer strategies that would enhance myocardial oxygenation and to study whether it improves the survival and engraftment of transplanted stem cells.

Myocardial oxygenation under conditions of ischemia-reperfusion

The most common cause of MI is narrowing of the epicardial blood vessels due to atheromatous plaques. Plaque ruptures, with subsequent exposure of the basement membrane, result in platelet aggregation, thrombus formation, fibrin accumulation, and hemorrhage into the plaque, followed by varying degrees of vasospasm. This can result in partial or complete occlusion of the vessel and subsequent myocardial ischemia. Total occlusion of the vessel for more than 4-6 hours results in irreversible myocardial necrosis, but reperfusion within this period can salvage the myocardium and reduce morbidity and mortality. Monitoring the myocardial tissue oxygenation may play a vital role in understanding the pathophysiology of oxygen changes that occur as a result of ischemia-reperfusion (I/R) injury in the heart, and it may serve as an important indicator to monitor the pathophysiology of several cardiovascular drugs that are administered following an acute myocardial infarction.

Recently, we have published several reports on pharmacological pre- and post-conditioning of rat hearts using various treatment remedy like verapamil (VER), HO-4038 (a verapamil derivative), sulfaphenazole (SPZ) and trimetazidine (TMZ) (Khan et al.; Khan et al. 2009b; Mohan et al. 2009). In all groups of hearts, the baseline myocardial pO_2 was about 20 mmHg, which dropped to 2-4 mmHg upon induction of regional ischemia by ligation of the left-anterior-descending (LAD) coronary artery (Figure 4A). After 30 min of ligation, blood flow was restored (reperfusion) by release of the ligation. Immediately upon reperfusion, there was a substantial hyperoxygenation in the untreated (saline only) I/R control group (Figure 4A). The myocardial oxygenation remained significantly elevated even after 48 h of reperfusion (Khan et al. 2010a). The hyperoxygenation may be due to contractile “stunning”, which is a reversible loss of contractility known to occur immediately upon reperfusion and last for several hours to days (Bolli and Marban 1999). During this time the myocardium receives adequate oxygen supply, but it does not fully utilize the oxygen because of depressed contractility (myocardial work) which is not immediately restored to preischemic levels. This condition may lead to a paradoxical hyperoxia due to the lack of oxygen utilization. Ambrosio et al have observed an overshoot of cardiac phosphocreatine concentration in rabbit hearts upon post-ischemic reflow (Ambrosio et al. 1987). This overshoot effect was attributed to a decrease in phosphocreatine utilization leading to an imbalance between supply and rate of utilization of the high energy phosphate metabolic reserve in the “stunned” heart (Ambrosio et al. 1987). Similarly, in yet another study the marked hyperoxygenation was attributed to decreased oxygen consumption due to NO-mediated inhibition of mitochondrial respiration (Zhao et al. 2005).

This reperfusion-induced hyperoxygenation was significantly attenuated in verapamil, HO-4038, and TMZ-treated groups when compared to untreated I/R controls (Figure 4A). The reduction in hyperoxygenation observed in the treated groups may be attributed to increase in oxygen demand due to improved recovery of cardiac function, as much as 3-fold increase in RPP (Rate pressure Product) in the HO-4038 group compared to control (Mohan et al. 2009). On the other hand, pretreatment of rats with SPZ showed a substantial hyperoxygenation at 1-h reperfusion (Figure 4A), which was attributed to enhanced NO levels that may promote increased blood flow upon reperfusion (Khan et al. 2009b). The marked hyperoxygenation could also occur as a result of decreased oxygen consumption due to NO-mediated inhibition of mitochondrial respiration (Khan et al. 2009b). In a recent study we observed that TMZ ameliorated the I/R-induced oxygen overshoot (Khan et al. 2010a). The absence of hyperoxygenation (Figure 4A) at reperfusion in the TMZ-treated hearts could be associated with an increased recovery of contractility and attenuation of myocardial injury. Also, the hyperoxygenation at reperfusion might indicate a decrease in oxygen consumption by the injured tissue.

Effect of post-administration of hyperoxygenation on stem-cell treatment for myocardial infarction

Hyperbaric oxygenation (HBO) is a safe, clinically-viable treatment that has been used as a primary therapy in patients with decompression sickness, arterial gas embolism and carbon monoxide poisoning (Tibbles and Edelsberg 1996). It is also used as an adjuvant therapy to promote wound healing (Thackham et al. 2008), and for the treatment of various conditions, including ischemic injury (Yogarathnam et al. 2006). HBO involves inhalation of 100% oxygen under greater-than-one atmospheric absolute (ATA) pressure. Such doses of oxygen have a number of beneficial biochemical, cellular, and physiologic effects (Yogarathnam et al. 2007). HBO, administered at the onset of reperfusion in an open-chest rabbit model of myocardial ischemia–reperfusion injury, showed a significant reduction in infarct size (Sterling et al. 1993). More recent studies have also shown that HBO attenuates myocardial injury *via* nitric oxide signaling (Yogarathnam et al. 2007), improves cardiac function in patients with acute myocardial infarction (Dekleva et al. 2004), and helps mobilization of stem cells by stimulating nitric oxide synthesis (Thom et al. 2006) and enhancing CXCR4 and VEGFR-2 in humans (Thom et al. 2006). However, until recently, the efficacy of HBO as an adjuvant to cell therapy had not yet been studied.

We used periods of HBO exposure in combination with myocardial stem-cell therapy to treat rat hearts subjected to MI. HBO exposure (100% O₂, 2 ATA, 90 min daily for 2 weeks) started three days after cell transplantation to allow the animals recover from surgical trauma. Administration of HBO, in conjunction with MSCs, increased tissue oxygenation in the infarct heart as measured by EPR spectroscopy (Khan et al. 2009a). *In vivo* myocardial pO₂ measurements were obtained at baseline and at the end of 2 weeks in all groups, including the rats that were receiving stem-cell therapy applied to the infarct heart. The mean baseline pO₂ in healthy hearts prior to simulated MI was 21.25±2.0 mmHg (Figure 4B). The myocardial pO₂ in hearts receiving MSC transplants was significantly improved when compared to untreated MI hearts (9.8±2.3 vs 3.8±1.2 mmHg, respectively). An additional group given adjuvant HBO therapy after receiving MSC transplants to the MI heart had further significant (p<0.05) enhancement in myocardial oxygenation when compared to the MSC-alone treated group (16.2±2.2 vs 9.8±2.3, respectively) (Figure 4B). The pO₂ data indicated that HBO therapy further increased myocardial oxygenation values to near-normal in infarct rat hearts treated with transplanted stem cells. Most importantly, this study showed that HBO treatment lead to increased stem-cell engraftment and improved cardiac function (Khan et al. 2009a).

EPR spectroscopy is a unique technique by which we can monitor non-invasively the myocardial oxygenation in the infarct heart. Our lab has previously published long-term monitoring of myocardial oxygenation, for up to 3 months, following OxySpin implantation in a healthy mouse heart (Khan et al. 2008). As mentioned earlier in our previous reports with skeletal myoblast transplantation (Khan et al. 2007), there was a marginal improvement in myocardial oxygenation. However, the combined treatment of stem cells along with hyperbaric oxygen (HBO) therapy showed a significant enhancement in myocardial oxygenation and functional recovery in the infarct heart (Khan et al. 2009a). The oximetry studies clearly demonstrated the role of oxygen in myocardial infarction and its importance in the treatment of cardiac repair by stem-cell therapy.

Overall, we have demonstrated the use of EPR oximetry for noninvasive monitoring of local tissue oxygenation in hearts treated with stem cells. The results clearly show that tissue oxygenation is an important factor for the survival and engraftment of stem cells in the heart.

Acknowledgments

We acknowledge the grant support from NIH (EB006153, EB004031) and from AHA (SDG 0930181N).

References

- Agbulut O, Vandervelde S, Al Attar N, Larghero J, Ghostine S, Leobon B, Robidel E, Borsani P, Le Lorc'h M, Bissery A, Chomienne C, Bruneval P, Marolleau JP, Vilquin JT, Hagege A, Samuel JL, Menasche P. Comparison of human skeletal myoblasts and bone marrow-derived CD133+ progenitors for the repair of infarcted myocardium. *Journal of the American College of Cardiology*. 2004; 44(2):458–463. [PubMed: 15261948]
- Ahmad R, Kuppusamy P. Theory, Instrumentation, and Applications of Electron Paramagnetic Resonance Oximetry. *Chemical Reviews*. 2010; 110(5):3212–3236. [PubMed: 20218670]
- Ambrosio G, Jacobus WE, Bergman CA, Weisman HF, Becker LC. Preserved high energy phosphate metabolic reserve in globally “stunned” hearts despite reduction of basal ATP content and contractility. *Journal of Molecular and Cellular Cardiology*. 1987; 19(10):953–964. [PubMed: 3437454]
- Arnesen H, Lunde K, Aakhus S, Forfang K. Cell therapy in myocardial infarction. *Lancet*. 2007; 369(9580):2142–2143. [PubMed: 17604783]
- Bolli R, Marban E. Molecular and cellular mechanisms of myocardial stunning. *Physiological Reviews*. 1999; 79(2):609–634. [PubMed: 10221990]
- Chacko SM, Khan M, Kuppusamy ML, Pandian RP, Varadharaj S, Selvendiran K, Bratasz A, Rivera BK, Kuppusamy P. Myocardial oxygenation and functional recovery in infarct rat hearts transplanted with mesenchymal stem cells. *American Journal of Physiology (Heart and Circulatory Physiology)*. 2009; 296(5):H1263–1273. [PubMed: 19286938]
- Chen XD, Dusevich V, Feng JQ, Manolagas SC, Jilka RL. Extracellular matrix made by bone marrow cells facilitates expansion of marrow-derived mesenchymal progenitor cells and prevents their differentiation into osteoblasts. *Journal of Bone and Minerals Research*. 2007; 22(12):1943–1956.
- Chiu RC, Zibaitis A, Kao RL. Cellular cardiomyoplasty: myocardial regeneration with satellite cell implantation. *The Annals of Thoracic Surgery*. 1995; 60(1):12–18. [PubMed: 7598572]
- D'Ippolito G, Diabira S, Howard GA, Roos BA, Schiller PC. Low oxygen tension inhibits osteogenic differentiation and enhances stemness of human MIAMI cells. *Bone*. 2006; 39(3):513–522. [PubMed: 16616713]
- Dekleva M, Neskovic A, Vlahovic A, Putnikovic B, Beleslin B, Ostojic M. Adjunctive effect of hyperbaric oxygen treatment after thrombolysis on left ventricular function in patients with acute myocardial infarction. *American Heart Journal*. 2004; 148(4):E14. [PubMed: 15459609]

- Djouad F, Delorme B, Maurice M, Bony C, Apparailly F, Louis-Plence P, Canovas F, Charbord P, Noel D, Jorgensen C. Microenvironmental changes during differentiation of mesenchymal stem cells towards chondrocytes. *Arthritis Research & Therapy*. 2007; 9(2):R33. [PubMed: 17391539]
- Egeland T, Brinchmann JE. Cell quality in the ASTAMI study. *European Heart Journal*. 2007a; 28(17):2172. author reply 2173-2174. [PubMed: 17675309]
- Egeland T, Brinchmann JE. The REPAIR-AMI and ASTAMI trials: cell isolation procedures. *European Heart Journal*. 2007b; 28(17):2174–2175. author reply 2175. [PubMed: 17673446]
- Ezashi T, Das P, Roberts RM. Low O₂ tensions and the prevention of differentiation of hES cells. *Proceedings of the National Academy of Sciences of the United States of America*. 2005; 102(13): 4783–4788. [PubMed: 15772165]
- Fehrer C, Brunauer R, Laschober G, Unterluggauer H, Reitingner S, Kloss F, Gully C, Gassner R, Lepperdinger G. Reduced oxygen tension attenuates differentiation capacity of human mesenchymal stem cells and prolongs their lifespan. *Aging cell*. 2007; 6(6):745–757. [PubMed: 17925003]
- Fink T, Abildtrup L, Fogd K, Abdallah BM, Kassem M, Ebbesen P, Zachar V. Induction of adipocyte-like phenotype in human mesenchymal stem cells by hypoxia. *Stem Cells*. 2004; 22(7):1346–1355. [PubMed: 15579652]
- Ghostine S, Carrion C, Souza LC, Richard P, Bruneval P, Vilquin JT, Pouzet B, Schwartz K, Menasche P, Hagege AA. Long-term efficacy of myoblast transplantation on regional structure and function after myocardial infarction. *Circulation*. 2002; 106(12 Suppl 1):I131–136. [PubMed: 12354722]
- Heidt MC, Sedding D, Stracke SK, Stadlbauer T, Boening A, Vogt PR, Schonburg M. Measurement of myocardial oxygen tension: a valid and sensitive method in the investigation of transmyocardial laser revascularization in an acute ischemia model. *The Thoracic and Cardiovascular Surgeon*. 2009; 57(2):79–84. [PubMed: 19241308]
- Kamihata H, Matsubara H, Nishiue T, Fujiyama S, Tsutsumi Y, Ozono R, Masaki H, Mori Y, Iba O, Tateishi E, Kosaki A, Shintani S, Murohara T, Imaizumi T, Iwasaka T. Implantation of bone marrow mononuclear cells into ischemic myocardium enhances collateral perfusion and regional function via side supply of angioblasts, angiogenic ligands, and cytokines. *Circulation*. 2001; 104(9):1046–1052. [PubMed: 11524400]
- Khan M, Kutala VK, Vikram DS, Wisel S, Chacko SM, Kuppasamy ML, Mohan IK, Zweier JL, Kwiatkowski P, Kuppasamy P. Skeletal myoblasts transplanted in the ischemic myocardium enhance in situ oxygenation and recovery of contractile function. *American Journal of Physiology (Heart and Circulatory Physiology)*. 2007; 293(4):H2129–2139. [PubMed: 17660391]
- Khan M, Kutala VK, Wisel S, Chacko SM, Kuppasamy ML, Kwiatkowski P, Kuppasamy P. Measurement of oxygenation at the site of stem cell therapy in a murine model of myocardial infarction. *Advances in Experimental Medicine and Biology*. 2008:61445–52.
- Khan M, Meduru S, Mohan IK, Kuppasamy ML, Wisel S, Kulkarni A, Rivera BK, Hamlin RL, Kuppasamy P. Hyperbaric oxygenation enhances transplanted cell graft and functional recovery in the infarct heart. *Journal of Molecular and Cellular Cardiology*. 2009a; 47(2):275–287. [PubMed: 19376124]
- Khan M, Meduru S, Mostafa M, Khan S, Hideg K, Kuppasamy P. Trimetazidine, administered at the onset of reperfusion, ameliorates myocardial dysfunction and injury by activation of p38 mitogen-activated protein kinase and Akt signaling. *The Journal of Pharmacology and Experimental Therapeutics*. 2010a; 333(2):421–429. [PubMed: 20167841]
- Khan M, Mohan IK, Kutala VK, Kotha SR, Parinandi NL, Hamlin RL, Kuppasamy P. Sulfaphenazole protects heart against ischemia-reperfusion injury and cardiac dysfunction by overexpression of iNOS, leading to enhancement of nitric oxide bioavailability and tissue oxygenation. *Antioxidants & Redox Signaling*. 2009b; 11(4):725–738. [PubMed: 18855521]
- Lane, N. *Oxygen: The Molecule that made the World*. Oxford University Press; Oxford: 2002.
- Lennon DP, Edmison JM, Caplan AI. Cultivation of rat marrow-derived mesenchymal stem cells in reduced oxygen tension: effects on in vitro and in vivo osteochondrogenesis. *Journal of Cellular Physiology*. 2001; 187(3):345–355. [PubMed: 11319758]

- Ma T, Grayson WL, Frohlich M, Vunjak-Novakovic G. Hypoxia and stem cell-based engineering of mesenchymal tissues. *Biotechnology Progress*. 2009; 25(1):32–42. [PubMed: 19198002]
- Martin-Rendon E, Brunskill S, Doree C, Hyde C, Watt S, Mathur A, Stanworth S. Stem cell treatment for acute myocardial infarction. *Cochrane Database of Systematic Reviews (Online)*. 2008a; (4):CD006536.
- Martin-Rendon E, Brunskill SJ, Hyde CJ, Stanworth SJ, Mathur A, Watt SM. Autologous bone marrow stem cells to treat acute myocardial infarction: a systematic review. *European Heart Journal*. 2008b; 29(15):1807–1818. [PubMed: 18523058]
- McConnell PI, del Rio CL, Jacoby DB, Pavlicova M, Kwiatkowski P, Zawadzka A, Dinsmore JH, Astra L, Wisel S, Michler RE. Correlation of autologous skeletal myoblast survival with changes in left ventricular remodeling in dilated ischemic heart failure. *The Journal of Thoracic and Cardiovascular Surgery*. 2005; 130(4):1001. [PubMed: 16214511]
- Mik EG, Ince C, Eerbeek O, Heinen A, Stap J, Hooibrink B, Schumacher CA, Balestra GM, Johannes T, Beek JF, Nieuwenhuis AF, van Horssen P, Spaan JA, Zuurbier CJ. Mitochondrial oxygen tension within the heart. *Journal of Molecular and Cellular Cardiology*. 2009; 46(6):943–951. [PubMed: 19232352]
- Mohan IK, Khan M, Wisel S, Selvendiran K, Sridhar A, Carnes CA, Bognar B, Kalai T, Hideg K, Kuppusamy P. Cardioprotection by HO-4038, a novel verapamil derivative, targeted against ischemia and reperfusion-mediated acute myocardial infarction. *American Journal of Physiology (Heart and Circulatory Physiology)*. 2009; 296(1):H140–151. [PubMed: 18978191]
- Orlic D, Kajstura J, Chimenti S, Bodine DM, Leri A, Anversa P. Transplanted adult bone marrow cells repair myocardial infarcts in mice. *Annals of the New York Academy of Sciences*. 2001a:938221–229. discussion 229–230.
- Orlic D, Kajstura J, Chimenti S, Jakoniuk I, Anderson SM, Li B, Pickel J, McKay R, Nadal-Ginard B, Bodine DM, Leri A, Anversa P. Bone marrow cells regenerate infarcted myocardium. *Nature*. 2001b; 410(6829):701–705. [PubMed: 11287958]
- Pagani FD, DerSimonian H, Zawadzka A, Wetzel K, Edge AS, Jacoby DB, Dinsmore JH, Wright S, Aretz TH, Eisen HJ, Aaronson KD. Autologous skeletal myoblasts transplanted to ischemia-damaged myocardium in humans. Histological analysis of cell survival and differentiation. *Journal of the American College of Cardiology*. 2003; 41(5):879–888. [PubMed: 12628737]
- Pandian RP, Dang V, Manoharan PT, Zweier JL, Kuppusamy P. Effect of nitrogen dioxide on the EPR property of lithium octa-n-butoxy 2,3-naphthalocyanine (LiNc-BuO) microcrystals. *Journal of Magnetic Resonance*. 2006; 181(1):154–161. [PubMed: 16690337]
- Pandian RP, Parinandi NL, Ilangovan G, Zweier JL, Kuppusamy P. Novel particulate spin probe for targeted determination of oxygen in cells and tissues. *Free Radical Biology & Medicine*. 2003; 35(9):1138–1148. [PubMed: 14572616]
- Potier E, Ferreira E, Andriamanalijaona R, Pujol JP, Oudina K, Logeart-Avramoglou D, Petite H. Hypoxia affects mesenchymal stromal cell osteogenic differentiation and angiogenic factor expression. *Bone*. 2007; 40(4):1078–1087. [PubMed: 17276151]
- Ren H, Cao Y, Zhao Q, Li J, Zhou C, Liao L, Jia M, Zhao Q, Cai H, Han ZC, Yang R, Chen G, Zhao RC. Proliferation and differentiation of bone marrow stromal cells under hypoxic conditions. *Biochemical and Biophysical Research Communications*. 2006; 347(1):12–21. [PubMed: 16814746]
- Retuerto MA, Schalch P, Patejunas G, Carbray J, Liu N, Esser K, Crystal RG, Rosengart TK. Angiogenic pretreatment improves the efficacy of cellular cardiomyoplasty performed with fetal cardiomyocyte implantation. *The Journal of Thoracic and Cardiovascular Surgery*. 2004; 127(4):1041–1049. discussion 1049–1051. [PubMed: 15052201]
- Rochefort GY, Delorme B, Lopez A, Heralut O, Bonnet P, Charbord P, Eder V, Domenech J. Multipotential mesenchymal stem cells are mobilized into peripheral blood by hypoxia. *Stem Cells*. 2006; 24(10):2202–2208. [PubMed: 16778152]
- Salim A, Giaccia AJ, Longaker MT. Stem cell differentiation. *Nature Biotechnology*. 2004; 22(7):804–805. author reply 805–806.
- Seeger FH, Tonn T, Krzossok N, Zeiher AM, Dimmeler S. Cell isolation procedures matter: a comparison of different isolation protocols of bone marrow mononuclear cells used for cell

- therapy in patients with acute myocardial infarction. *European Heart Journal*. 2007; 28(6):766–772. [PubMed: 17298974]
- Silverman HS, Wei S, Haigney MC, Ocampo CJ, Stern MD. Myocyte adaptation to chronic hypoxia and development of tolerance to subsequent acute severe hypoxia. *Circulation Research*. 1997; 80(5):699–707. [PubMed: 9130451]
- Sterling DL, Thornton JD, Swafford A, Gottlieb SF, Bishop SP, Stanley AW, Downey JM. Hyperbaric oxygen limits infarct size in ischemic rabbit myocardium in vivo. *Circulation*. 1993; 88(4 Pt 1): 1931–1936. [PubMed: 8403338]
- Strauer BE, Brehm M, Zeus T, Kosterling M, Hernandez A, Sorg RV, Kogler G, Wernet P. Repair of infarcted myocardium by autologous intracoronary mononuclear bone marrow cell transplantation in humans. *Circulation*. 2002; 106(15):1913–1918. [PubMed: 12370212]
- Swartz HM, Clarkson RB. The measurement of oxygen in vivo using EPR techniques. *Physics in Medicine and Biology*. 1998; 43(7):1957–1975. [PubMed: 9703059]
- Thackham JA, McElwain DL, Long RJ. The use of hyperbaric oxygen therapy to treat chronic wounds: A review. *Wound Repair and Regeneration*. 2008; 16(3):321–330. [PubMed: 18471250]
- Thom SR, Bhopale VM, Velazquez OC, Goldstein LJ, Thom LH, Buerk DG. Stem cell mobilization by hyperbaric oxygen. *American Journal of Physiology (Heart and Circulatory Physiology)*. 2006; 290(4):H1378–1386. [PubMed: 16299259]
- Thom T, Haase N, Rosamond W, Howard VJ, Rumsfeld J, Manolio T, Zheng ZJ, Flegal K, O'Donnell C, Kittner S, Lloyd-Jones D, Goff DC Jr, Hong Y, Adams R, Friday G, Furie K, Gorelick P, Kissela B, Marler J, Meigs J, Roger V, Sidney S, Sorlie P, Steinberger J, Wasserthiel-Smolter S, Wilson M, Wolf P. Heart disease and stroke statistics--2006 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation*. 2006; 113(6):e85–151. [PubMed: 16407573]
- Tibbles PM, Edelsberg JS. Hyperbaric-oxygen therapy. *The New England Journal of Medicine*. 1996; 334(25):1642–1648. [PubMed: 8628361]
- Tse HF, Kwong YL, Chan JK, Lo G, Ho CL, Lau CP. Angiogenesis in ischaemic myocardium by intramyocardial autologous bone marrow mononuclear cell implantation. *Lancet*. 2003; 361(9351):47–49. [PubMed: 12517468]
- van Beem RT, Hirsch A, Lommerse IM, Zwaginga JJ, Noort WA, Biemond BJ, Piek JJ, van der Schoot CE, Voermans C. Recovery and functional activity of mononuclear bone marrow and peripheral blood cells after different cell isolation protocols used in clinical trials for cell therapy after acute myocardial infarction. *Eurointervention*. 2008; 4(1):133–138. [PubMed: 19112790]
- Vogt S, Troitzsch D, Spath S, Portig I, Moosdorf R. Direct measurement of myocardial oxygen tension and high energy phosphate content under varying ventilatory conditions in rabbits. *Biomedizinische Technik*. 2009; 54(4):179–186. [PubMed: 19807283]
- Wang DW, Fermor B, Gimble JM, Awad HA, Guilak F. Influence of oxygen on the proliferation and metabolism of adipose derived adult stem cells. *Journal of Cellular Physiology*. 2005; 204(1):184–191. [PubMed: 15754341]
- Wisel S, Chacko SM, Kuppusamy ML, Pandian RP, Khan M, Kutala VK, Burry RW, Sun B, Kwiatkowski P, Kuppusamy P. Labeling of skeletal myoblasts with a novel oxygen-sensing spin probe for noninvasive monitoring of in situ oxygenation and cell therapy in heart. *American Journal of Physiology (Heart and Circulatory Physiology)*. 2007; 292(3):H1254–1261. [PubMed: 17142337]
- Yogarathnam JZ, Laden G, Guvendik L, Cowen M, Cale A, Griffin S. Can hyperbaric oxygen be used as adjunctive heart failure therapy through the induction of endogenous heat shock proteins? *Advances in Therapy*. 2007; 24(1):106–118. [PubMed: 17526467]
- Yogarathnam JZ, Laden G, Madden LA, Seymour AM, Guvendik L, Cowen M, Greenman J, Cale A, Griffin S. Hyperbaric oxygen: a new drug in myocardial revascularization and protection? *Cardiovascular Revascularization Medicine*. 2006; 7(3):146–154. [PubMed: 16945821]
- Zhao X, He G, Chen YR, Pandian RP, Kuppusamy P, Zweier JL. Endothelium-derived nitric oxide regulates postischemic myocardial oxygenation and oxygen consumption by modulation of mitochondrial electron transport. *Circulation*. 2005; 111(22):2966–2972. [PubMed: 15939832]




Figure 1. Representative EPR spectra obtained from OxySpin in the heart. EPR spectra of a suspension of OxySpin in PBS equilibrated with room air (**A**) and 100% nitrogen (**B**). EPR spectra obtained from an infarcted mouse heart, *in vivo*, 1 day (**C**) or 2 weeks (**D**) after implantation of OxySpin-labeled skeletal myoblast (SM) cells. The peak-peak width (w) of the spectrum is a measure of oxygen concentration. The measured pO_2 values are noted on the respective spectrum.




Figure 2. Long-term monitoring of myocardial pO₂ values in beating hearts. The measurements were performed repeatedly for 16 weeks using *in vivo* EPR oximetry in murine hearts implanted with microcrystals of LiNc-BuO (OxySpin) probe in the mid-ventricular region without LAD coronary artery ligation. Data represent mean±SD;(n=6). The results show the feasibility of pO₂ measurements for more than 3 months in the beating hearts of mice.




Figure 3. Myocardial pO₂ in the infarcted heart at the site of cell transplantation. Myocardial pO₂ values were measured repeatedly for 4 weeks using *in vivo* EPR oximetry in murine hearts transplanted with OxySpin-labeled SM cells. **(A)** Tissue pO₂ at 4 weeks after treatment with SM cells (MI+SM) was significantly higher when compared to untreated (MI) hearts. **(B)** The time-course values of myocardial pO₂ measured from infarcted hearts (MI), and infarcted hearts treated with SM cells (MI+SM) are shown. Values are expressed as mean \pm SD; (n=7). *p<0.05 versus MI group.




Figure 4. Values of pO₂ in rat hearts under conditions of ischemia-reperfusion and stem-cell therapy for myocardial infarction. **(A)** Effect of pretreatment of rats with verapamil (VER), HO-4038, SPZ, and TMZ on myocardial pO₂ during regional ischemia followed by reperfusion. Myocardial tissue pO₂ values were measured by EPR oximetry at 30 min of ischemia, followed by at 1 h of reperfusion. *p<0.05 versus Saline group; n=6. **(B)** Measurement of myocardial pO₂ in rat hearts transplanted with mesenchymal stem cells (MSC) at 2 weeks. The results show an increase in myocardial oxygenation levels in hearts treated with MSC and further enhancement in pO₂ was observed in hearts treated in conjunction with HBO and MSCs. The “baseline” data were obtained from hearts before

induction of ischemia. Data represent mean \pm SD; n=6. *p<0.05 *versus* Saline group.
**p<0.05 *versus* MSC group.